
Kansas: Most Common Causes of Death

- Total cardiovascular diseases (principally ischemic heart disease and stroke) are the most common cause of death in Kansas, all cancers are the second most common cause, chronic obstructive pulmonary disease is fourth, and diabetes is sixth.
- In 1995, 72% of all deaths in Kansas were due to the four most common chronic disease causes of death.
- The death rate for chronic obstructive pulmonary disease was higher in Kansas than in the United States; the death rates for total cardiovascular diseases, all cancers, and diabetes were lower.

Causes of Death, Kansas Compared With United States, 1995*




^{*}All data are age adjusted, 1970 total U.S. population.

^{**}Total cardiovascular disease death rates include rates of death due to ischemic heart disease (120.9 per 100,000 in Kansas and 135.2 per 100,000 in the United States) and rates of death due to stroke (41.0 per 100,000 in Kansas and 42.5 per 100,000 in the United States).


Kansas: Cardiovascular Diseases

- Total cardiovascular diseases are the most common cause of death in Kansas, accounting for 39% of all deaths.
- Ischemic heart disease accounted for 49% of all cardiovascular disease deaths in Kansas in 1995; 4,827 people in Kansas died of ischemic heart disease.
- In 1995, 1,811 people in Kansas died of stroke.


Kansas: Total Cardiovascular Disease Death Rates



Kansas: Ischemic Heart Disease Death Rates




Kansas: Stroke Death Rates





Note: All data are age adjusted, 1970 total U.S. population.


Kansas: Cancer

- Cancer accounted for 25% of all deaths in Kansas in 1995;
 5,283 people in Kansas died of cancer.
- In Kansas in 1995, 1,427 people died of lung cancer, 568 people died of colorectal cancer, and 433 women died of breast cancer.
- The American Cancer Society estimates that 1,600 new cases of lung cancer, 1,300 new cases of colorectal cancer, and 1,600 new cases of breast cancer will be diagnosed in Kansas in 1997.










Note: All data are age adjusted, 1970 total U.S. population.

Kansas


Kansas: Diabetes

- In 1994, 81,525 adults in Kansas had diagnosed diabetes.
- Diabetes was the underlying cause of 577 deaths in Kansas in 1995.
- In 1993, diabetes was the most common contributing cause of 192 new cases of end-stage kidney disease in Kansas.
- Diabetes is believed to be underreported on death certificates, both as a condition and as a cause of death.



Note: All data are age adjusted, 1970 total U.S. population.

Kansas: Risk Factors




 ${}^\star\textsc{Ever}$ smoked at least 100 cigarettes and now smoke every day or some days.

Kansas: Percentage of Adults Who Reported No Leisure-Time Physical Activity* 100 By Sex, Race, and Hispanic Origin, 1996 BRFSS 80 57.3 Percent 46.5 36.2 34.8 40 35.3 34.3 20 0 Total Male Female White Black Other Hispanic

*No exercise, recreation, or physical activity (other than regular job duties) during the previous month.

^{**}Too few numbers to analyze.



*Body mass index \geq 27.8 kg/m² for men and \geq 27.3 kg/m² for women.

Kansas: Percentage of High School Students Who Reported Cigarette Smoking

By Sex, 1995 YRBSS

YRBSS data not available

Kansas: Percentage of High School Students Not Enrolled in Physical Education Classes

By Sex, 1995 YRBSS

YRBSS data not available


Note: All BRFSS data are provisional and age adjusted, 1970 total U.S. population.

Kansas

^{**}Too few numbers to analyze.

^{**}Too few numbers to analyze.

Kansas: Risk Factors




Kansas: Percentage of High School Students Who Reported Eating Fewer Than Five Servings of Fruits and Vegetables on the Day Preceding the Survey By Sex, 1995 YRBSS

YRBSS data not available

11.6

Hispanic

Kansas: Preventive Services



Note: All BRFSS data are provisional and age adjusted, 1970 total U.S. population.

Kansas